

Genome-Scale CRISPR-Cas9 Knockout Screening in Human Cells

Ophir Shalem,^{1,2*} Neville E. Sanjana,^{1,2*} Ella Hartenian,¹ Xi Shi,^{1,3} David A. Scott,^{1,2} Tarjei Mikkelsen,¹ Dirk Heckl,⁴ Benjamin L. Ebert,⁴ David E. Root,¹ John G. Doench,¹ Feng Zhang^{1,2†}

¹Broad Institute of MIT and Harvard, 7 Cambridge Center, Cambridge, MA 02142, USA. ²McGovern Institute for Brain Research, Department of Brain and Cognitive Sciences, Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, USA. ³Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT, 7 Cambridge Center, Cambridge, MA 02142, USA. ⁴Division of Hematology, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA 02115, USA.

*These authors contributed equally to this work.

†Corresponding author. E-mail: zhang@broadinstitute.org

The simplicity of programming the CRISPR-associated nuclease Cas9 to modify specific genomic loci suggests a new way to interrogate gene function on a genome-wide scale. We show that lentiviral delivery of a genome-scale CRISPR-Cas9 knockout (GeCKO) library targeting 18,080 genes with 64,751 unique guide sequences enables both negative and positive selection screening in human cells. First, we used the GeCKO library to identify genes essential for cell viability in cancer and pluripotent stem cells. Next, in a melanoma model, we screened for genes whose loss is involved in resistance to vemurafenib, a therapeutic that inhibits mutant protein kinase BRAF. Our highest-ranking candidates include previously validated genes *NF1* and *MED12* as well as novel hits *NF2*, *CUL3*, *TADA2B*, and *TADA1*. We observe a high level of consistency between independent guide RNAs targeting the same gene and a high rate of hit confirmation, demonstrating the promise of genome-scale screening with Cas9.

A major goal since the completion of the Human Genome Project is the functional characterization of all annotated genetic elements in normal biological processes and disease (1). Genome-scale loss-of-function screens have provided a wealth of information in diverse model systems (2–5). In mammalian cells, RNA interference (RNAi) is the predominant method for genome-wide loss-of-function screening (2, 3), but its utility is limited by the inherent incompleteness of protein depletion by RNAi and confounding off-target effects (6, 7).

The RNA-guided CRISPR (clustered regularly interspaced short palindromic repeats)-associated nuclease Cas9 provides an effective means of introducing targeted loss-of-function mutations at specific sites in the genome (8, 9). Cas9 can be programmed to induce DNA double strand breaks (DSBs) at specific genomic loci (8, 9) through a synthetic single guide RNA (sgRNA) (10), which when targeted to coding regions of genes can create frame shift indel mutations that result in a loss-of-function allele. Because the targeting specificity of Cas9 is conferred by short guide sequences, which can be easily generated at large scale by array-based oligonucleotide library synthesis (11), we sought to explore the potential of Cas9 for pooled genome-scale functional screening.

Lentiviral vectors are commonly used for delivery of pooled short hairpin RNAs (shRNAs) in RNAi since they can be easily titrated to control transgene copy number, and are stably maintained as genomic integrants during subsequent cell replication (2, 12, 13). Therefore we designed a single lentiviral vector to deliver Cas9, a sgRNA, and a puromycin selection marker into target cells (lentiCRISPR) (Fig. 1A). The ability to simultaneously deliver Cas9 and sgRNA through a single vector enables application to any cell type of interest, without the need to first generate cell lines that express Cas9.

To determine the efficacy of gene knockout by lentiCRISPR trans-

duction, we tested six sgRNAs targeting enhanced green fluorescent protein (EGFP) in a HEK293T cell line containing a single copy of EGFP (fig. S1). After transduction at a low multiplicity of infection (MOI = 0.3) followed by selection with puromycin, lentiCRISPRs abolished EGFP fluorescence in $93 \pm 8\%$ (mean \pm s.d.) of cells after 11 days (Fig. 1B). Deep sequencing of the EGFP locus revealed a $92 \pm 9\%$ indel frequency ($n \geq 10^4$ sequencing reads per condition) (fig. S2). In contrast, transduction of cells with lentiviral vectors expressing EGFP-targeting shRNA led to incomplete knockdown of EGFP fluorescence (Fig. 1C).

Given the high efficacy of gene knockout by lentiCRISPR, we tested the feasibility of conducting genome-scale CRISPR-Cas9 knockout (GeCKO) screening with a pooled lentiCRISPR library. We designed a library of sgRNAs targeting 5' constitutive exons (Fig. 2A) of 18,080 genes in the human genome with an average coverage of 3–4 sgRNAs per gene (table S1), and each target site was selected to minimize off-target modification (14) (supplementary discussion).

To test the efficacy of the full GeCKO library at achieving knock-out of endogenous gene targets, we conducted a negative selection screen by profiling the depletion of sgRNAs targeting essential survival genes (Fig. 2A). We transduced the human melanoma cell line A375 and the human stem cell line HUES62 with the GeCKO library at a MOI of 0.3. As expected, deep sequencing (figs. S3 and S4) 14 days post-transduction revealed a significant reduction in the diversity of sgRNAs in the surviving A375 and HUES62 cells (Fig. 2, B and C) (Wilcoxon rank sum test, $P < 10^{-10}$ for both cell types). Gene set enrichment analysis (GSEA) (15) indicated that most of the depleted sgRNAs targeted essential genes such as ribosomal structural constituents (Fig. 2, D and E, and tables S2 and S3). The overlap in highly depleted genes and functional gene categories between the two cell lines (fig. S5) indicates that GeCKO can identify essential genes and that enrichment analysis of depleted sgRNAs can pinpoint gene targets in negative selection screens. To test the efficacy of GeCKO for positive selection, we sought to identify gene knockouts that result in resistance to the BRAF protein kinase inhibitor vemurafenib (PLX) in melanoma (16) (Fig. 3A). Exposure to PLX resulted in growth arrest of transduced A375 cells, which harbor the V600E gain-of-function BRAF mutation (17) (Fig. 3B), therefore enabling the enrichment of a small group of cells that were rendered drug-resistant by Cas9:sgRNA-mediated modification. After 14 days of PLX treatment, the sgRNA distribution was significantly different when compared with vehicle-treated cells (Fig. 3C) (Wilcoxon rank-sum test, $P < 10^{-10}$) and clustered separately from all other conditions (Fig. 3D and fig. S6).

For a subset of genes, we found enrichment of multiple sgRNAs that target each gene after 14 days of PLX treatment (Fig. 3E), suggesting that loss of these particular genes contributes to PLX resistance. We used the RNAi Gene Enrichment Ranking (RIGER) algorithm to rank

screening hits by the consistent enrichment among multiple sgRNAs targeting the same gene (Fig. 3F and table S4) (12). Our highest ranking genes included previously reported candidates *NF1* and *MED12* (18, 19) and also several genes not previously implicated in PLX resistance, including neurofibromin 2 (*NF2*), Cullin 3 E3 ligase (*CUL3*), and members of the STAGA histone acetyltransferase complex (*TADA1* and *TADA2B*). These candidates yield new testable hypotheses regarding PLX resistance mechanisms (supplementary discussion). For example, *NF1* and *NF2*, although unrelated in sequence, are each mutated in similar but distinct forms of neurofibromatosis (20). In addition, epigenetic dysregulation resulting from mutations in the mechanistically related STAGA and Mediator complexes (21) may have a role in acquired drug resistance. All of these hits were also identified through a second independent transduction (figs. S7 and S8, and tables S5 and S6).

A similar screen to identify PLX drug resistance in A375 cells was previously conducted using a pooled library of 90,000 shRNAs (19). To compare the efficacy and reliability of genome-scale shRNA screening with GeCKO, we used several methods to evaluate the degree of consistency among the sgRNAs or shRNAs targeting the top candidate genes. First, we calculated the aggregate *P* value distribution for the top 100 hits using either RIGER (Fig. 4A) or RSA (fig. S9) scoring. Lower *P* values for the GeCKO versus shRNA screen indicate better scoring consistency among sgRNAs. Second, for the top 10 RIGER hit genes, $78 \pm 27\%$ of sgRNAs targeting each gene ranked among the top 5% of enriched sgRNAs, whereas $20 \pm 12\%$ of shRNAs targeting each gene ranked among the top 5% of enriched shRNAs (Fig. 4B).

We validated top ranking genes from the GeCKO screen individually using 3-5 sgRNAs (Fig. 4, C to E, and figs. S10 and S11). For *NF2*, we found that 4/5 sgRNAs resulted in >98% allele modification 7 days post-transduction, and all 5 sgRNAs showed >99% allele modification 14 days post-transduction (Fig. 4C). We compared sgRNA and shRNA-mediated protein depletion and PLX resistance using Western blot (Fig. 4D) and cell growth assays (Fig. 4E). Interestingly, while all five sgRNAs conferred resistance to PLX, only the best shRNA achieved sufficient knockdown to increase PLX resistance (Fig. 4E), suggesting that even low levels of *NF2* are sufficient to retain sensitivity to PLX. Additionally, sgRNAs targeting *NF1*, *MED12*, *CUL3*, *TADA1*, and *TADA2B* led to a decrease in protein expression and increased resistance to PLX (figs. S10 and S11). Deep sequencing confirmed a high rate of mutagenesis at targeted loci (figs. S12 and S13), with a small subset of off-target sites exhibiting indels (figs. S14 to S16), which may be alleviated using an offset nicking approach (22, 23) that was recently shown to reduce off-target modifications (22).

GeCKO screening provides a mechanistically distinct method to RNAi for systematic perturbation of gene function. Whereas RNAi reduces protein expression by targeting RNA, GeCKO introduces loss-of-function mutations into genomic DNA. While some indel mutations are expected to maintain the reading frame, homozygous knockout yields high screening sensitivity, which is especially important in cases where incomplete knockdown retains gene function. In addition, RNAi is limited to transcripts, whereas Cas9:sgRNAs can target elements across the entire genome, including promoters, enhancers, introns, and intergenic regions. Furthermore, catalytically inactive mutants of Cas9 can be tethered to different functional domains (23–27) to broaden the repertoire of perturbation modalities, including genome-scale gain-of-function screening using Cas9-activators and epigenetic modifiers. In the GeCKO screens presented here, the efficiency of complete knockout, the consistency of distinct sgRNAs, and the validation rate for top screen hits demonstrate the potential of Cas9:sgRNA-based technology to transform functional genomics.

References and Notes

1. E. S. Lander, Initial impact of the sequencing of the human genome.

- Nature* **470**, 187–197 (2011). [doi:10.1038/nature09792](https://doi.org/10.1038/nature09792) [Medline](#)
- K. Berns, E. M. Hijmans, J. Mullenders, T. R. Brummelkamp, A. Velds, M. Heimerikx, R. M. Kerkhoven, M. Madiredjo, W. Nijkamp, B. Weigelt, R. Agami, W. Ge, G. Cavet, P. S. Linsley, R. L. Beijersbergen, R. Bernards, A large-scale RNAi screen in human cells identifies new components of the p53 pathway. *Nature* **428**, 431–437 (2004). [doi:10.1038/nature02371](https://doi.org/10.1038/nature02371) [Medline](#)
 - M. Boutros, A. A. Kiger, S. Armknecht, K. Kerr, M. Hild, B. Koch, S. A. Haas, R. Paro, N. Perrimon; Heidelberg Fly Array Consortium, Genome-wide RNAi analysis of growth and viability in *Drosophila* cells. *Science* **303**, 832–835 (2004). [doi:10.1126/science.1091266](https://doi.org/10.1126/science.1091266) [Medline](#)
 - R. Rad, L. Rad, W. Wang, J. Cadinanos, G. Vassiliou, S. Rice, L. S. Campos, K. Yusa, R. Banerjee, M. A. Li, J. de la Rosa, A. Strong, D. Lu, P. Ellis, N. Conte, F. T. Yang, P. Liu, A. Bradley, PiggyBac transposon mutagenesis: A tool for cancer gene discovery in mice. *Science* **330**, 1104–1107 (2010). [doi:10.1126/science.1193004](https://doi.org/10.1126/science.1193004) [Medline](#)
 - J. E. Carette, C. P. Guimaraes, M. Varadarajan, A. S. Park, I. Wuethrich, A. Godarova, M. Kotecki, B. H. Cochran, E. Spooner, H. L. Ploegh, T. R. Brummelkamp, Haploid genetic screens in human cells identify host factors used by pathogens. *Science* **326**, 1231–1235 (2009). [doi:10.1126/science.1178955](https://doi.org/10.1126/science.1178955) [Medline](#)
 - A. L. Jackson, J. Burchard, J. Schelter, B. N. Chau, M. Cleary, L. Lim, P. S. Linsley, Widespread siRNA “off-target” transcript silencing mediated by seed region sequence complementarity. *RNA* **12**, 1179–1187 (2006). [doi:10.1261/rna.25706](https://doi.org/10.1261/rna.25706) [Medline](#)
 - C. J. Echeverri, P. A. Beachy, B. Baum, M. Boutros, F. Buchholz, S. K. Chanda, J. Downward, J. Ellenberg, A. G. Fraser, N. Hacohen, W. C. Hahn, A. L. Jackson, A. Kiger, P. S. Linsley, L. Lum, Y. Ma, B. Mathey-Prévôt, D. E. Root, D. M. Sabatini, J. Taipale, N. Perrimon, R. Bernards, Minimizing the risk of reporting false positives in large-scale RNAi screens. *Nat. Methods* **3**, 777–779 (2006). [doi:10.1038/nmeth1006-777](https://doi.org/10.1038/nmeth1006-777) [Medline](#)
 - L. Cong, F. A. Ran, D. Cox, S. Lin, R. Barretto, N. Habib, P. D. Hsu, X. Wu, W. Jiang, L. A. Marraffini, F. Zhang, Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**, 819–823 (2013). [doi:10.1126/science.1231143](https://doi.org/10.1126/science.1231143) [Medline](#)
 - P. Mali, L. Yang, K. M. Esvelt, J. Aach, M. Guell, J. E. DiCarlo, J. E. Norville, G. M. Church, RNA-guided human genome engineering via Cas9. *Science* **339**, 823–826 (2013). [doi:10.1126/science.1232033](https://doi.org/10.1126/science.1232033) [Medline](#)
 - M. Jinek, K. Chylinski, I. Fonfara, M. Hauer, J. A. Doudna, E. Charpentier, A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816–821 (2012). [doi:10.1126/science.1225829](https://doi.org/10.1126/science.1225829) [Medline](#)
 - A. P. Blanchard, L. Hood, Sequence to array: Probing the genome’s secrets. *Nat. Biotechnol.* **14**, 1649 (1996). [doi:10.1038/nbt1296-1649](https://doi.org/10.1038/nbt1296-1649) [Medline](#)
 - B. Luo, H. W. Cheung, A. Subramanian, T. Sharifnia, M. Okamoto, X. Yang, G. Hinkle, J. S. Boehm, R. Beroukhi, B. A. Weir, C. Mermel, D. A. Barbie, T. Awad, X. Zhou, T. Nguyen, B. Piqani, C. Li, T. R. Golub, M. Meyerson, N. Hacohen, W. C. Hahn, E. S. Lander, D. M. Sabatini, D. E. Root, Highly parallel identification of essential genes in cancer cells. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 20380–20385 (2008). [doi:10.1073/pnas.0810485105](https://doi.org/10.1073/pnas.0810485105) [Medline](#)
 - P. J. Paddison, J. M. Silva, D. S. Conklin, M. Schlabach, M. Li, S. Aruleba, V. Balija, A. O’Shaughnessy, L. Gnoj, K. Scobie, K. Chang, T. Westbrook, M. Cleary, R. Sachidanandam, W. R. McCombie, S. J. Elledge, G. J. Hannon, A resource for large-scale RNA-interference-based screens in mammals. *Nature* **428**, 427–431 (2004). [doi:10.1038/nature02370](https://doi.org/10.1038/nature02370) [Medline](#)
 - P. D. Hsu, D. A. Scott, J. A. Weinstein, F. A. Ran, S. Konermann, V. Agarwala, Y. Li, E. J. Fine, X. Wu, O. Shalem, T. J. Cradick, L. A. Marraffini, G. Bao, F. Zhang, DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat. Biotechnol.* **31**, 827–832 (2013). [doi:10.1038/nbt.2647](https://doi.org/10.1038/nbt.2647) [Medline](#)
 - A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette, A. Paulovich, S. L. Pomeroy, T. R. Golub, E.

- S. Lander, J. P. Mesirov, Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 15545–15550 (2005). [doi:10.1073/pnas.0506580102](https://doi.org/10.1073/pnas.0506580102) [Medline](#)
16. K. T. Flaherty, I. Puzanov, K. B. Kim, A. Ribas, G. A. McArthur, J. A. Sosman, P. J. O'Dwyer, R. J. Lee, J. F. Grippo, K. Nolop, P. B. Chapman, Inhibition of mutated, activated BRAF in metastatic melanoma. *N. Engl. J. Med.* **363**, 809–819 (2010). [doi:10.1056/NEJMoa1002011](https://doi.org/10.1056/NEJMoa1002011) [Medline](#)
 17. H. Davies, G. R. Bignell, C. Cox, P. Stephens, S. Edkins, S. Clegg, J. Teague, H. Woffendin, M. J. Garnett, W. Bottomley, N. Davis, E. Dicks, R. Ewing, Y. Floyd, K. Gray, S. Hall, R. Hawes, J. Hughes, V. Kosmidou, A. Menzies, C. Mould, A. Parker, C. Stevens, S. Watt, S. Hooper, R. Wilson, H. Jayatilake, B. A. Gusterson, C. Cooper, J. Shipley, D. Hargrave, K. Pritchard-Jones, N. Maitland, G. Chenevix-Trench, G. J. Riggins, D. D. Bigner, G. Palmieri, A. Cossu, A. Flanagan, A. Nicholson, J. W. Ho, S. Y. Leung, S. T. Yuen, B. L. Weber, H. F. Seigler, T. L. Darrow, H. Paterson, R. Marais, C. J. Marshall, R. Wooster, M. R. Stratton, P. A. Futreal, Mutations of the BRAF gene in human cancer. *Nature* **417**, 949–954 (2002). [doi:10.1038/nature00766](https://doi.org/10.1038/nature00766) [Medline](#)
 18. S. Huang, M. Hölzel, T. Knijnenburg, A. Schlicker, P. Roepman, U. McDermott, M. Garnett, W. Grenrum, C. Sun, A. Prahallad, F. H. Groenendijk, L. Mittempergher, W. Nijkamp, J. Neeffjes, R. Salazar, P. Ten Dijke, H. Uramoto, F. Tanaka, R. L. Beijersbergen, L. F. Wessels, R. Bernards, MED12 controls the response to multiple cancer drugs through regulation of TGF- β receptor signaling. *Cell* **151**, 937–950 (2012). [doi:10.1016/j.cell.2012.10.035](https://doi.org/10.1016/j.cell.2012.10.035) [Medline](#)
 19. S. R. Whittaker, J. P. Theurillat, E. Van Allen, N. Wagle, J. Hsiao, G. S. Cowley, D. Schadendorf, D. E. Root, L. A. Garraway, A genome-scale RNA interference screen implicates NF1 loss in resistance to RAF inhibition. *Cancer Discovery* **3**, 350–362 (2013). [doi:10.1158/2159-8290.CD-12-0470](https://doi.org/10.1158/2159-8290.CD-12-0470) [Medline](#)
 20. A. L. Lin, D. H. Gutmann, Advances in the treatment of neurofibromatosis-associated tumours. *Nat. Rev. Clin. Oncol.* **10**, 616–624 (2013). [doi:10.1038/nrclinonc.2013.144](https://doi.org/10.1038/nrclinonc.2013.144) [Medline](#)
 21. X. Liu, M. Vorontchikhina, Y. L. Wang, F. Faiola, E. Martinez, STAGA recruits Mediator to the MYC oncoprotein to stimulate transcription and cell proliferation. *Mol. Cell. Biol.* **28**, 108–121 (2008). [doi:10.1128/MCB.01402-07](https://doi.org/10.1128/MCB.01402-07) [Medline](#)
 22. F. A. Ran, P. D. Hsu, C. Y. Lin, J. S. Gootenberg, S. Konermann, A. E. Trevino, D. A. Scott, A. Inoue, S. Matoba, Y. Zhang, F. Zhang, Double nicking by RNA-guided CRISPR Cas9 for enhanced genome editing specificity. *Cell* **154**, 1380–1389 (2013). [doi:10.1016/j.cell.2013.08.021](https://doi.org/10.1016/j.cell.2013.08.021) [Medline](#)
 23. P. Mali, J. Aach, P. B. Stranges, K. M. Esvelt, M. Moosburner, S. Kosuri, L. Yang, G. M. Church, CAS9 transcriptional activators for target specificity screening and paired nickases for cooperative genome engineering. *Nat. Biotechnol.* **31**, 833–838 (2013). [doi:10.1038/nbt.2675](https://doi.org/10.1038/nbt.2675) [Medline](#)
 24. L. A. Gilbert, M. H. Larson, L. Morsut, Z. Liu, G. A. Brar, S. E. Torres, N. Stern-Ginossar, O. Brandman, E. H. Whitehead, J. A. Doudna, W. A. Lim, J. S. Weissman, L. S. Qi, CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* **154**, 442–451 (2013). [doi:10.1016/j.cell.2013.06.044](https://doi.org/10.1016/j.cell.2013.06.044) [Medline](#)
 25. S. Konermann, M. D. Brigham, A. E. Trevino, P. D. Hsu, M. Heidenreich, L. Cong, R. J. Platt, D. A. Scott, G. M. Church, F. Zhang, Optical control of mammalian endogenous transcription and epigenetic states. *Nature* **500**, 472–476 (2013). [Medline](#)
 26. P. Perez-Pinera, D. D. Kocak, C. M. Vockley, A. F. Adler, A. M. Kabadi, L. R. Polstein, P. I. Thakore, K. A. Glass, D. G. Ousterout, K. W. Leong, F. Guilak, G. E. Crawford, T. E. Reddy, C. A. Gersbach, RNA-guided gene activation by CRISPR-Cas9-based transcription factors. *Nat. Methods* **10**, 973–976 (2013). [doi:10.1038/nmeth.2600](https://doi.org/10.1038/nmeth.2600) [Medline](#)
 27. M. L. Maeder, S. J. Linder, V. M. Cascio, Y. Fu, Q. H. Ho, J. K. Joung, CRISPR RNA-guided activation of endogenous human genes. *Nat. Methods* **10**, 977–979 (2013). [doi:10.1038/nmeth.2598](https://doi.org/10.1038/nmeth.2598) [Medline](#)

Acknowledgments: We thank G. Cowley, W. Harrington, J. Wright, E.

Hodis, S. Whittaker, J. Merkin, C. Burge, L. P. Club, and the entire Zhang laboratory for technical support and critical discussions. O.S. is a Klarman Fellow, N.S. is a Simons Fellow, D.A.S. is an NSF Fellow, and J.D. is a Merkin Institute Fellow. F.Z. is supported by an NIH Director's Pioneer Award (1DP1-MH100706), a NIH Transformative R01 grant (1R01-DK097768), the Keck, McKnight, Merkin, Vallee, Damon Runyon, Searle Scholars, Klingenstein, and Simons Foundations, Bob Metcalfe, and Jane Pauley. The authors have no conflicting financial interests. A patent application has been filed relating to this work, and the authors plan to make the reagents widely available to the academic community through Addgene and to provide software tools via the Zhang laboratory Web site (www.genome-engineering.org).

Supplementary Materials

www.sciencemag.org/cgi/content/full/science.1247005/DC1

Materials and Methods

Supplementary Text

Figs. S1 to S16

Tables S1 to S10

References

9 October 2013; accepted 2 December 2013

Published online 12 December 2013

10.1126/science.1247005




Fig. 1. Lentiviral delivery of Cas9 and sgRNA provides efficient depletion of target genes. (A) Lentiviral expression vector for Cas9 and sgRNA (lentiCRISPR). Puromycin selection marker (puro), psi packaging signal (psi+), rev response element (RRE), central polypurine tract (cPPT), elongation factor-1 α short promoter (EFS), 2A self-cleaving peptide (P2A), and posttranscriptional regulatory element (WPRE). (B) Distribution of fluorescence from 293T-EGFP cells transduced by EGFP-targeting lentiCRISPR (sgRNAs 1-6, outlined peaks) and Cas9-only (green-shaded peak) vectors, and non-fluorescent 293T cells (gray shaded peak). (C) Distribution of fluorescence from 293T-EGFP cells transduced by EGFP-targeting shRNA (shRNAs 1-4, outlined peaks) and control shRNA (green-shaded peak) vectors, and non-fluorescent 293T cells (gray shaded peak).




Fig. 2. GeCKO library design and application for genome-scale negative selection screening. (A) Design of sgRNA library for genome-scale knockout of coding sequences in human cells (supplementary discussion). (B and C) Cumulative frequency of sgRNAs 3 and 14 days post transduction in A375 and hES cells respectively. Shift in the 14 day curve represents the depletion in a subset of sgRNAs. (D and E) Five most significantly depleted gene sets in A375 cells (nominal $P < 10^{-5}$, FDR-corrected $q < 10^{-5}$) and HUES62 cells (nominal $P < 10^{-5}$, FDR-corrected $q < 10^{-3}$) identified by Gene Set Enrichment Analysis (GSEA) (15).




Fig. 3. GeCKO screen in A375 melanoma cells reveals genes whose loss confers vemurafenib (PLX) resistance.

(A) Timeline of PLX resistance screen in A375 melanoma cells. (B) Growth of A375 cells when treated with DMSO or PLX over 14 days. (C) Boxplot showing the distribution of sgRNA frequencies at different time points, with and without PLX treatment (vehicle = DMSO). The box extends from the first to the third quartile with the whiskers denoting 1.5 times the interquartile range. Enrichment of specific sgRNAs: 7 days of PLX treatment, 1 sgRNA greater than 10-fold enrichment; 14 days of PLX treatment, 379 and 49 sgRNAs greater than 10-fold and 100-fold enrichment respectively. (D) Rank correlation of normalized sgRNA read count between biological replicates and treatment conditions. (E) Scatterplot showing enrichment of specific sgRNAs after PLX treatment. (F) Identification of top candidate genes using the RNAi Gene Enrichment Ranking (RIGER) P value analysis.




Fig. 4. Comparison of GeCKO and shRNA screens and validation of neurofibromin 2 (*NF2*). (A) RIGER P values for the top 100 hits from GeCKO and shRNA (19) screens for genes whose loss results in PLX resistance. Analysis using the Redundant siRNA Activity (RSA) algorithm shows a similar trend (fig. S9). (B) For the top 10 RIGER hits, the percent of unique sgRNAs (top) or shRNAs (bottom) targeting each gene that are in top 5% of all enriched sgRNAs or shRNAs. (C) Deep sequencing analysis of lentiCRISPR-mediated indel at the *NF2* locus. (D) A375 cells transduced with *NF2*-targeting lentiCRISPR and shRNA vectors both show a decrease in *NF2* protein levels. (E) Dose response curves for A375 cells transduced with individual *NF2*-targeting lentiCRISPR or shRNA vectors. Controls were EGFP-targeting lentiCRISPR or null hairpin shRNA vectors. Cells transduced with *NF2*-targeting lentiCRISPRs show a significant increase ($F_{1,8} = 30.3$, $P < 0.001$, $n = 4$ replicates) in the half maximal effective concentration (EC_{50}) whereas cells transduced with *NF2*-targeting shRNA vectors do not ($F_{1,8} = 0.47$, $P = 0.51$, $n = 4$ replicates).